



Vitamin Content and Storage Studies of Cookies Produced from Wheat, Almond and Carrot Flour Blends

Ahure Dinnah¹, Mulak Desmond Guyih^{1*} and Mike O. Eke²

¹*Department of Food Science and Technology, Federal University of Agriculture, Makurdi, Nigeria.*

²*Department of Chemistry, Faculty of Science, CEFTER, Benue State University, Makurdi, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author MDG conducted the experiment and collected the data and analyzed them. Authors AD and MOE contributed in supervision of the experiment, paper formatting, helped to check the English, grammar and also processing process during submission to the journal. All authors read and approved the final manuscript.

Article Information

DOI:10.9734/AFSJ/2020/v15i330154

Editor(s):

(1) Dr. PapiyaBigoniya, RKDF University, India.

(2) Dr. Nelson Pérez Guerra, University of Vigo, Spain.

Reviewers:

(1)Oluwatoyin A. Oladeji,OlusegunAgagu University of Science and Technology (OAUSTECH), Nigeria.

(2) Tamer El Sisy, Agriculture Research Center, Egypt.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/55670>

Original Research Article

Received 01 February 2020

Accepted 04 April 2020

Published 15 May 2020

ABSTRACT

The purpose of this study was to produce cookies from wheat, almond and carrot flour blend, evaluate the vitamin content and storage parameters. Wheat, almond and carrot flour were blended in the ratio: 100:0:0, 90:10:0, 90:0:10, 80:15:5, 70:20:10 and were labeled A, B, C, D and E respectively to produce cookies. The control sample A was without almond and carrot flour. The cookies produced were analysed for vitamin content and were stored for 7weeks at relative humidity corresponding to wet and dry season condition (70% and 30% respectively). The cookies were then analysed for pH, moisture and fungi content in an interval of every 2 weeks using standard methods, at the end of the storage, the sensory attributes and vitamin content of the cookies were analysed. The vitamin content range: from 341.53 to 653.27 µg/100 g for vitamin A, 1.523 to 2.450 mg/g for vitamin B1, 0.65 to 0.92 mg/g for vitamin B2, 3.12 to 3.52 mg/g for vitamin B3 and 2.093 to 3.007 mg/g for vitamin C. All cookies samples were generally accepted by sensory panelist before storage and at the end of storage time. At the end of storage, pH value ranged from 5.5 to 7.8 for wet season condition cookies and from 5.5 to 5.7 for dry season condition cookies.

*Corresponding author: Email: mulakdesmond@gmail.com;

The moisture content ranged from 4.5 to 6.17% for wet season condition cookies and 1.33 to 1.63% for dry season condition cookies. The vitamin A content after storage ranged from 341.53 to 653.23 IU/100 g for wet season condition cookies and 336.61 to 653.01 IU/100 g for dry season condition cookies, while vitamin C ranged from 2.093 to 3.007 mg/g and 2.11 to 3.01 mg/g for wet and dry season condition cookies respectively. 1CFU of fungi was identified for each sample of cookie. The study provides evidence that wheat, almond and carrot are suitable for cookies production and variation of storage conditions did not cause spoilage of cookies.

Keywords: Vitamin; relative humidity; sensory attributes.

1. INTRODUCTION

Bakery cookies are very popular, ready to eat, convenient, less expensive and also important product in human diet. There are usually eaten with tea and used as weaning food for infants. It is also used as a snack in school for the school going children who are often underweight. It may be used as a nutrient supplement during emergency situation [1]. Cookie is also known as an excellent vehicle for incorporation of different nutritionally rich ingredients, thus making it a useful tool in meeting the nutritional requirements of the increasing population.

Wheat ranks first among the cereals used for baking in Nigeria and is associated with growth and survival of the people of the country [2]. More than 60% of the total daily requirements of protein and calories are met through wheat; wheat provides 360 kilo calories [3]. It contributes 68-75% of the total food intake in the daily diet and provides 75% of the total protein requirements [4]. It is a staple food, consumed worldwide in the form of bread and biscuits etc. [4]. The edible portion of almonds (*Prunus amygdalus*) is its nuts, which are commonly known as almonds or badam and it is a popular, nutritious food. Almond seeds contain 46% fat, 21% protein, 25% carbohydrate and appreciable amount of mineral elements [5]. The majority of lipids in almond are monounsaturated (~67%) and polyunsaturated (~25%) fatty acids (MUFA and PUFA, respectively). Previous studies indicate that the MUFAs from almond seeds may reduce total cholesterol and low-density lipoproteins (LDL, "bad cholesterol") while maintaining healthy high-density lipoproteins (HDL, "good cholesterol") levels [6]. Carrots are a root vegetable also known as *Daucus carota*, it is one of the top ten foods that food consumers identify as having "health benefits beyond basic nutrition". This reputation comes with good reason, as carrots are rich in carotenoids which are used to make vitamin A during digestion, and

vitamins B, C, D and E [7]. They're also high in folic acid, fiber, and minerals like Potassium (K) and Sodium (Na). Sweet and succulent carrot are notably rich in anti-oxidants, vitamins and dietary fiber; however, they provide only 41 calories per 100 g, negligible amount of fat and no cholesterol. They are exceptionally rich source of carotenes and vitamin-A. 100 g fresh carrot contains 8285 mcg of beta-carotene and 16706 IU of vitamin A [8].

Adequate nutrition during infancy and early childhood is fundamental to the development of each child's full human potential. It is well recognized that the period from birth to two years of age is a "critical window" for the promotion of optimal growth, health and behavioral development. Longitudinal studies have consistently shown that this is the peak age for growth faltering, deficiencies of certain micronutrients, and common childhood illnesses such as diarrhea [9]. Food fortification a process of adding micronutrients to processed foods can be used to tackle this problem of nutritional deficiency. In many situations, this strategy can lead to relatively rapid improvements in the micronutrient status of a population, and at a very reasonable cost, especially if advantage can be taken of existing technology and local distribution networks [10]. Since the benefits are potentially large, food fortification can be cost-effective in public health intervention [11]. Composting cookies with wheat, almond and carrot flour is therefore expected to increase the micronutrient content of cookies.

2. MATERIALS AND METHODS

2.1 Procurement of Raw Materials

Carrots, almond seeds, margarine, sugar, eggs, baking powder and wheat were purchased from the modern market Makurdi, Benue State of Nigeria. Procured materials were taken to the Chemistry department, Benue State University laboratory for processing.

2.2 Preparation of Flour

Almond seeds was sorted/cleaned, washed and then oven dried at 60°C for 2 hours and then ground into flour using laboratory grinders (M/S Sujata: New Delhi India). The almond flour was sieved through a 0.5 mm size mesh. Carrots were sorted, cleaned, washed and cut into smaller sizes of 1 cm cubes and then blanched at 70°C for 5 minutes. The carrot cubes was oven dried at 50°C for 24 hours and then milled into flour. The carrot flour was then sieved through a 0.5 mm size mesh. Wheat was wet milled into flour according to [12].

2.3 Preparation of Composite Flour Blends

Composite flour was prepared from wheat flour fortified with almond seeds flour and carrot flour at different levels (Table 1). Cookies were prepared from supplemented flour with modification of the method described in [12] as indicated in Table 2.

2.4 Method

The creaming method was used for the preparation of dough where vegetable fat and sugar were creamed together using a Kenwood mixer (United Kingdom) at medium speed for two (2) min. After creaming, such ingredients as flour, baking powder and egg were added and mixed to form dough and properly mixed (Table 2). The dough was manually needed to ensure uniformity and then transferred to a clean tray and gently rolled using a roller. The dough sheath was cut

into round shapes using a cutter. Shaped dough pieces were placed into a greased pan and baked in an oven at 180°C for 40min. The baked cookies were placed on a cooling rack for 30 min to cool before packaging [12].

2.4.1 Vitamin analysis (Vitamin A)

Vitamin A was determined by the method described by [13]. Approximately 1 g of the sample was mixed with 30 ml of absolute alcohol and 3 ml of 5% KOH solution was added to it and was boiled for 30 min under reflux. After washing with distilled water, vitamin A was extracted with 150 ml of diethyl ether. The extract was evaporated to dryness at low temperature and then dissolved in 10 mL of isopropyl alcohol. Exactly 1 mL of standard vitamin A solution was prepared and that of the dissolved extract was transferred to separate cuvettes and their respective absorbance are read in a spectrophotometer at 325 nm with a reagent blank at zero.

2.4.2 Determination of thiamine (Vitamin B1)

The spectrophotometric method, described by Krishnaiah et al. [14] was used for determination of the B Vitamins. Exactly 5 g of each sample was homogenized with 50 ml of 1N ethanolic sodium hydroxide and the homogenate was filtered to obtain the filtrate to be used for the analysis. An aliquot (10 mL) of the filtrate was treated with equal volume of 0.1N K₂Cr₂O₇ solution in a flask. Standard thiamine solution was prepared and diluted to a chosen concentration (0.5).

$$\text{Conc. of vit A in sample} = \frac{\text{Abs of sample}}{\text{Abs of std}} \times \text{Conc. of STD} \quad (1)$$

Table1. Composition of flour blends

Samples	Wheat: almond: carrot flour ratio
A	100:0:0
B	90:10:0
C	90:0:10
D	85:15:5
E	70:20:10

Table 2. Recipe for the formulation of cookies from wheat flour, almond seed and carrot flour blends

Component	Cookies composition
Flour (g)	100
Butter (g)	45
Sugar (g)	45
Egg (mL)	30 ml
Baking powder (g)	2

An aliquot of the standard thiamine solution was also treated with 10 mL of the dichromate solution ($K_2Cr_2O_7$) in a separate flask while a reagent blank was set up by treating 10 mL of the ethanolic sodium hydroxide with the potassium dichromate solution. The absorbance of the sample and the standard solutions was measured in a spectrophotometer at a wavelength of 360 nm, with the reagent blank to be used to calibrate the instrument at zero. The thiamine content was calculated using the formula:

$$\text{Thiamine } \frac{mg}{100} = \frac{100}{W} \times \frac{Au}{As} \times \frac{C}{1} \times \frac{Vf}{Va} \times D \quad (2)$$

Where: W = Weight of sample analyzed, Au = Absorbance of sample, As = Absorbance of standard solution, C = Concentration (mg/ml) of standard solution, Vf = Total volume of filtrate, Va = Volume of filtrate analyzed, D = Dilution factor where applicable.

2.4.3 Determination of riboflavin (Vitamin B2)

Approximately 1 g of sample was weighed into a conical flask and was dissolved with 100 mL of deionized water. This was shaken thoroughly and heated for 5 min and allowed to cool and then filtered. The filtrate was poured into cuvettes and their respective wavelengths for the vitamins set to read the absorbance using spectrophotometer.

Vitamin B1 = 261nm, Vitamin B2 = 242 nm

$$\text{Vitamin conc. (mg/\%)} = A \times Df \times \text{Vol. of cuvette} \quad (3)$$

Where: A = Absorbance, E = Extinction coefficient = 25 for B1 and B2, Df = Dilution factor

2.4.4 Determination of niacin (Vitamin B3)

A measured weight (5 g) of each sample was treated with 50 ml of 1N sulphuric acid (H_2SO_4 solution) and was shaken for 30 min. The mixture was treated further with 3 drops of aqueous ammonia and filtered. The filtrate (extract) was used for the analysis. Standard niacin (nicotinic acid) solution was prepared and diluted as desired. 10 mL portion of the standard solution, sample extract and 10 mL of the acid solution (treated with a drop of ammonia) was dispensed into separate flasks to serve as standard, the sample and reagent blank respectively. Each of them was treated with 5 ml of normal potassium cyanide solution and acidified with 5 ml of 0.02N

H_2SO_4 solution; its absorbance was read in a spectrophotometer at a wavelength of 470 nm. The reagent blank was used to calibrate the instrument at zero. Niacin content was calculated using the formula [14];

$$\text{Niacine } \frac{mg}{100} = \frac{100}{W} \times \frac{Au}{As} \times \frac{C}{1} \times \frac{Vf}{Va} \times D \quad (4)$$

Where; W - Weight of sample analyzed, Au = Absorbance of sample, As = Absorbance of standard solution, C = Concentration (mg/ml) of standard solution, Vf = Total volume of filtrate, Va = Volume of filtrate analyzed, D = Dilution factor where applicable, C = Concentration of standard solution.

2.4.5 Determination of ascorbic acid (Vitamin C)

The method described by Krishnaiah et al. [14] was used. Exactly 10g of the sample was extracted with 50ml EDTA/TCA (50g in 50mL of water) extracting Solution for 1 hour and filtered through a Whatman filter paper into a 50mL volumetric flask and made up to the mark with the extracting solution. Twenty (20mL) of the extract was pipetted into a 250 mL conical flask and 10mL of 30% KI was added and also 50mL of distilled water added. This was followed by 2 mL of 1% starch indicator. It was titrated against 0.01 mL $CuSO_4$ solution to a dark end point.

$$\text{Vit. C } \frac{mg}{100g} = 0.88 \times \frac{100}{5} \times \frac{Vf}{20} \times \frac{T}{1} \quad (5)$$

Where: Vf = Volume of extract, T = Sample titre – blank titre.

2.4.6 Sensory evaluation

Sensory evaluation of cookies was carried out using 30 panelist comprising MSc/PhD students and staff of the Food Science and Technology course, CEFTER BSU Makurdi. Cookies were made with same quantity as commercially sold cookies. Panelist were required to evaluate the aroma, appearance, taste, mouth feel and overall acceptability of the cookies using a 9-point Hedonic scale with 1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike slightly, 5 = neither like nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much, 9 = like extremely [15].

2.4.7 Storage stability study

The cookies were packaged in a LDPE (Low Density Polyethylene) packaging material and stored at different relative humidity (relative

humidity corresponding to the dry and wet season for Makurdi that is; 75% and 30%) for 7 weeks at ambient temperature. After an interval of 14 days, samples were analyzed for pH, moisture content, microbial growth (mold and yeast) and sensory evaluation. The samples were later analyzed for vitamin A, C content at the end of the storage period. Moisture content of cookies samples was determined by standard method described by AOAC [16].

2.4.8 Microbial Analysis

Potato dextrose agar (PDA) was used for yeast and mold. Serial dilutions were made for each sample and 1 ml of the appropriate dilution poured, 10^{-5} and 10^{-6} serial dilution were used for pour plate in triplicate on selective media. Culture media was incubated at 37°C temperature for 7 days. The molds growth from the samples was purely cultured and isolated/identified. Developed colonies were expressed as colony forming units per gram (cfu/g) of sample [16]. Vitamin A was analysed by the method described by AOAC [17] while vitamin C was determined by the method described by AOAC [16].

2.5 Statistical Analysis

Statistical package for social science (SPSS) V21 computer software was used to analyze the data. Mean and standard deviation was calculated where appropriate. Analysis of variance (one way ANOVA) was used to determine the treatment that were different from the others in the various parameters tested; differences was considered at 95% ($p < 0.05$) significant level and 99% ($p < 0.01$) significant level where mentioned. The Dorcan multiple test was used to separate the means.

3. RESULTS AND DISCUSSION

3.1 Vitamin Content of Cookies

Results for the vitamin content of cookies are shown on Table 3. There was a constant increase in the vitamin content of all the cookie samples. The control sample had the lowest amount of vitamin in each case. This increase in vitamin is as a result of increase fortification with almond and carrot flour which is higher in micronutrients than wheat flour. The significant increase in this vitamin A from 341.53 $\mu\text{g}/100\text{g}$ to 653.27 $\mu\text{g}/100\text{g}$ and vitamin B₁ from 1.523 to 2.450 mg/g with increase in almond and carrot

flour suggests that vitamin A and vitamin B₁ are concentrated in almond and carrot flour than wheat flour. The vitamin B₁ values obtained were higher than that of the control sample and are much higher than the 0.130 to 0.423 mg/100 g reported by Okafor and Ugwu [18] for ready to eat snacks made from blends of breadfruit, cashew nut and coconut. There was also an excess of the recommended minimum daily intake of 0.2 to 1.0 mg according to Belitz [19]. The vitamin B₂ increased from 0.65 mg/100g in the control sample to 0.92 mg/100 g in sample E, values where in range with the recommended daily intake of 0.3 to 1.6 mg [20]. This equally took similar pattern as in Vitamins A and B₁ and can be attributed to the rich deposit of the B vitamins in almond and carrot. The values are much higher than the 0.157 to 0.477 mg/100 g reported by Okafor and Ugwu [18]. The B vitamins are needed for carbohydrate and protein metabolism, and are essential for growth, well structuring and functioning of the cells. There was significant ($p < 0.05$) difference in Vitamin C content of the samples with Sample E excelling with 3.01 mg/100 g while sample A was the least with 2.09 mg/100 g. The values compared lower with the value 4.77 mg/100 g reported by Ifediba and Obasi [21]. Vitamin C is a strong water soluble antioxidant that helps the body develop resistance against infectious agents and scavenges harmful free radicals. The vitamin A and C content of the cookie samples were not affected by storage conditions or storage time of 7 weeks; the values did not show any level of increase from the initial vitamin values Table 4. Vitamins are mostly aromatic shaped chemical compounds and degrade with time due to exposure to various environmental conditions e.g. vitamin A is susceptible to degradation by sun rays, other harsh environmental conditions like temperature, radiation exposure. Vitamins play vital functions in the body: they are needed for formation of hormones, cell growth, needed for proper functioning of the immune system, enhances clear vision etc.

3.2 Sensory Attributes of Cookies

Results of the sensory attributes of cookies are presented on Table 5. The values showed significant difference for its appearance and general acceptability while there were no significant difference in the texture, taste and aroma of the samples. All cookie samples were generally accepted by panelist, Sample A had the highest value of 7.74 closely followed by sample B (7.37). Sample A again had the highest

reading for appearance 7.89 and least in sample C (5.94). The statistical results reveal a positive score for all samples in the various parameters. These results show that cookies can be produced from wheat, almond and carrot flour blends at all substitution levels of 90:10:0, 90:0:10, 80:15:5, 70:20:10 (wheat, almond and carrot flour respectively). The results also showed that at continuous flour substitution levels, the sensory attributes start declining.

Utpal et al. [22], Bazilla et al. [23] reported the same trend of results for the sensory attributes of mushroom enriched biscuits and cookies produced from Sweet Potato- Maize Flour Blends respectively. The results for the sensory attributes of cookies stored at 75% and 30% relative humidity are shown in Table 6 and Table 7, the entire cookie samples were generally accepted by panelist for texture, taste, aroma, and appearance after 7 weeks of storage.

Table 3. Vitamin content of cookies

Sample	Vitamin A ($\mu\text{g}/100\text{g}$)	Vitamin B ₁ (mg/g)	Vitamin B ₂ (mg/g)	Vitamin B ₃ (mg/g)	Vitamin C (mg/g)
A	341.53 \pm 0.02 ^e	1.523 \pm 0.006 ^e	0.65 \pm 0.01 ^d	3.12 \pm 0.00 ^e	2.093 \pm 0.01 ^e
B	416.25 \pm 0.01 ^d	2.153 \pm 0.006 ^d	0.67 \pm 0.01 ^c	3.25 \pm 0.00 ^d	2.437 \pm 0.01 ^d
C	512.35 \pm 0.01 ^c	2.227 \pm 0.006 ^c	0.75 \pm 0.00 ^b	3.27 \pm 0.01 ^c	2.460 \pm 0.00 ^c
D	611.05 \pm 0.01 ^b	2.417 \pm 0.006 ^b	0.75 \pm 0.01 ^b	3.41 \pm 0.00 ^b	2.773 \pm 0.01 ^b
E	653.27 \pm 0.02 ^a	2.450 \pm 0.000 ^a	0.92 \pm 0.01 ^a	3.52 \pm 0.00 ^a	3.007 \pm 0.01 ^a

All data are means of 3 triplicates expressed on dry weight basis. Different superscripts between columns depict significant difference ($p \leq 0.05$)

Table 4. Increase level of Vitamin A and Vitamin C

Samples	Initial	(R.H 75%)	(R.H 30%)	Initial	(R.H 75%)	(R.H 30%)
Vitamin A ($\mu\text{g}/100\text{g}$)	week7	Vitamin C (mg/100g)week7		week7	week7	week7
A	341.53 \pm 0.02 ^e	337.00 \pm 0.03 ^e	336.61 \pm 0.02 ^e	2.093 \pm 0.01 ^e	2.10 \pm 0.02 ^e	2.11 \pm 0.01 ^e
B	416.25 \pm 0.01 ^d	418.03 \pm 0.12 ^d	418.23 \pm 0.12 ^d	2.437 \pm 0.01 ^d	2.39 \pm 0.05 ^d	2.40 \pm 0.05 ^d
C	512.35 \pm 0.01 ^c	511.23 \pm 0.09 ^c	510.96 \pm 0.01 ^c	2.460 \pm 0.00 ^c	2.50 \pm 0.23 ^c	2.51 \pm 0.06 ^c
D	611.05 \pm 0.01 ^b	610.02 \pm 0.07 ^b	611.87 \pm 0.03 ^b	2.773 \pm 0.01 ^b	2.77 \pm 0.01 ^b	2.66 \pm 0.12 ^b
E	653.27 \pm 0.02 ^a	652.01 \pm 0.13 ^a	653.01 \pm 0.01 ^a	3.007 \pm 0.01 ^a	3.00 \pm 0.21 ^a	3.01 \pm 0.17 ^a

All data are means of 3 triplicates expressed on dry weight basis. Different superscripts between columns depict significant difference ($p \leq 0.05$)

Table 5. Initial sensory attributes of cookies

Sample	Texture	Taste	Aroma	Appearance	General acceptability
A	7.50 \pm 1.29 ^a	7.56 \pm 1.20 ^a	7.11 \pm 1.49 ^a	7.89 \pm 0.96 ^a	7.74 \pm 1.10 ^a
B	7.50 \pm 1.50 ^a	7.44 \pm 1.50 ^a	6.94 \pm 1.59 ^a	7.50 \pm 1.10 ^{ab}	7.37 \pm 1.34 ^a
C	6.61 \pm 1.69 ^a	6.72 \pm 1.74 ^a	6.33 \pm 1.28 ^a	5.94 \pm 1.73 ^d	6.95 \pm 1.58 ^{ab}
D	7.00 \pm 1.53 ^a	7.33 \pm 1.37 ^a	6.94 \pm 1.30 ^a	6.72 \pm 1.27 ^{bcd}	7.05 \pm 1.27 ^a
E	7.06 \pm 1.59 ^a	7.17 \pm 1.58 ^a	6.67 \pm 1.78 ^a	6.17 \pm 1.72 ^{cd}	7.11 \pm 1.59 ^{ab}

Table 6. Sensory attributes of cookies stored at 75% relative humidity after 7 weeks

Samples	Texture	Taste	Aroma	Appearance	General acceptability
A	7.6 \pm 1.42 ^a	7.3 \pm 1.34 ^a	7.3 \pm 1.68 ^a	7.8 \pm 1.16 ^a	7.6 \pm 1.11 ^a
B	7.7 \pm 1.18 ^a	7.2 \pm 1.66 ^{ab}	7.0 \pm 1.34 ^{ab}	7.4 \pm 1.28 ^b	7.3 \pm 1.00 ^b
C	7.1 \pm 1.70 ^b	6.5 \pm 1.91 ^c	6.5 \pm 1.28 ^b	5.4 \pm 1.74 ^{bc}	6.8 \pm 1.25 ^{ab}
D	7.4 \pm 1.42 ^{ab}	7.2 \pm 1.47 ^{ab}	7.3 \pm 1.10 ^a	6.8 \pm 1.60 ^{ab}	6.9 \pm 1.30 ^{ab}
E	7.3 \pm 1.18 ^b	7.0 \pm 1.67 ^b	7.0 \pm 1.5 ^{ab}	6.1 \pm 1.70 ^c	7.3 \pm 1.90 ^b

Table 7. Sensory attributes of cookies stored at 30% relative humidity after 7weeks

Samples	Texture	Taste	Aroma	Appearance	General acceptability
A	7.5±1.14 ^b	7.3±1.32 ^a	7.2±1.61 ^a	7.8±1.23 ^a	7.65±1.43 ^a
B	7.8±1.23 ^a	7.15±1.45 ^a	7.0±1.27 ^a	7.3±1.45 ^b	7.2±1.12 ^b
C	6.8±1.26 ^{ab}	6.5±1.28 ^b	6.3±1.19 ^b	5.9±1.11 ^{ab}	6.8±1.56 ^{ab}
D	7.2±1.70 ^b	7.0±1.90 ^a	7.0±1.34 ^a	6.9±1.23 ^b	7.0±1.23 ^{ab}
E	7.4±1.54 ^b	6.9±1.34 ^a	7.0±1.56 ^a	6.3±1.45 ^{ab}	7.4±1.54 ^a

All data are means of 3 triplicates. Different superscripts between columns depict significant difference ($p \leq 0.05$)

3.3 Variation of Ph for Cookies Stored at 75% and 30% Relative Humidity Respectively

Cookies were stored for a period of 7 weeks at two different conditions that is; wet season condition (75% relative humidity) and dry season relative humidity (30% relative humidity). The pH values of cookies stored at 75% and 30% relative humidity were almost at the pH value of distilled water during the first 5 weeks of storage (Table 8). The cookies were within the acceptable pH range [24]. Usually, pH changes toward a high acidity or basicity depict spoilage by microorganisms (yeast and molds). The pH values of cookies decreased during the 7th week of storage towards the acid scale for cookies stored at wet season condition (7.8 to 5.5) and an acid scale value of 5.5 to 5.7 was recorded for cookies stored at dry season conditions. This difference in pH values can be attributed to the high percentage of moisture lost for cookies stored at 30% relative humidity (Table 8). Decreases in moisture content (water lost by vaporization) of food product increases the concentration of substances present in it the food product [24].

3.4 Variation of Moisture Content for Cookies Stored at 75% and 30% Relative Humidity Respectively

The results for cookies moisture content during storage are shown on Table 9: within the first five weeks of storage, there was a slight increase in the moisture content of cookies stored at wet season condition with a minimum value of 9.11% and a maximum value of 11.07%. These values later dropped at the end of the 7th week of storage 4.45% to 6.4% in sample A and D respectively. At the end of the storage time, a moisture content of 1.33 to 1.63% was recorded in sample A and E respectively. This decrease in moisture content for samples stored at different conditions signifies that, the moisture content of cookies during storage can be affected by the

environmental conditions of storage as well as the kind of packaging material used. These results showed that the cookies produced from wheat, almond and carrot flour blends were highly affected by humidity during the wet seasons (averagely 75% relative humidity) than dry seasons (averagely 30% relative humidity). Moisture (water) is very important for metabolic activities in living organisms. High moisture content in foodstuff provides the favourable environment for microbial activities. Nonetheless, the moisture content at the end of the 3th week slightly differ from the 9.05 to 10.25% initial moisture content of cookies produced from cassava, soybeans and mango flour blends reported by Chinma and Gernah [25]. The values at the 5th week of storage were in range with a 12.59 moisture reported by Lucretia et al. [26]. On the other hand, a decrease in moisture content depict less activity of spoilage organisms in food samples and the greater chance of the food product lasting for a longer period of time. The results of 4.5% to 6.4% moisture content of cookies stored at 75% relative humidity were similar to 5.0 to 6.1% moisture content of cookies produced from sweet potato- maize flour blends [27]. 2.73 to 3.01% moisture content was reported by Omachi and Yusufu [28] for soya flour carrot pomace biscuits which were similar to the moisture content of cookies stored at 30% relative humidity at the end of the studies.

3.5 Fungal Coliforms Identified Cookies Stored at 75% and 30% Relative Humidity

At the end of the storage period, 1cfu of fungi grew in each sample for both cookies stored at wet season condition and dry season condition (75 and 30% R.H respectively). For sample A *Saccharomyces spp* was identified for cookies stored at both seasonal conditions. These are flat, smooth, moist, glistening and creamy in colour, fungi of this group are good carbohydrate fermenters. They cause food spoilage of sugar rich foods such as cookies, syrup, juices etc.

Table 8. Variation of cookies pH during storage at 75% and 30% relative humidity

Samples	Wet season condition, pH at 75% R.H				Dry season condition, pH and 30% R.H)			
	Week 1	Week 3	Week 5	Week 7	Week 1	Week 3	Week 5	Week 7
A	7.21±0.01 ^d	7.21±0.01 ^d	7.21±0.01 ^d	5.5±0.0 ^c	7.19±0.00 ^a	7.19±0.0 ^a	7.19±0.00 ^a	5.5±0.0 ^a
B	7.16±0.01 ^e	7.17±0.01 ^e	7.17±0.00 ^e	6.6±0.0 ^b	7.18±0.0 ^b	7.17±0.0 ^b	7.18±0.01 ^b	5.6±0.0 ^a
C	7.27±0.01 ^b	7.28±0.01 ^b	7.28±0.00 ^b	5.5±0.0 ^c	7.18±0.0 ^b	7.17±0.0 ^b	7.18±0.00 ^b	5.5±0.0 ^a
D	7.25±0.01 ^c	7.25±0.01 ^c	7.26±0.01 ^c	7.8±0.0 ^a	7.15±0.01 ^c	7.15±0.0 ^c	7.15±0.01 ^c	5.7±0.0 ^a
E	7.30±0.01 ^a	7.30±0.01 ^a	7.30±0.01 ^a	6.67±0.0 ^b	7.19±0.00 ^a	7.19±0.0 ^a	7.19±0.00 ^a	5.6±0.0 ^a

All data are means of 3 replicates expressed on dry weight. Different superscripts between columns depict significant difference ($p \leq 0.05$)

Table 9. Variation of moisture content for cookies stored at 75% and 30% relative humidity respectively

Samples	(moisture at 75% and 30% R.H)							
	Week 1	Week 3	Week 5	Week 7	Week 1	Week 3	Week 5	Week 7
A	6.42±0.43 ^c	7.29±0.63 ^b	11.07±0.11 ^a	4.50±1.50 ^c	6.42±0.43 ^c	6.51±0.50 ^c	9.28±0.83 ^{bc}	1.33±0.58 ^b
B	7.43±0.45 ^{ab}	8.11±0.75 ^{ab}	10.98±0.06 ^a	6.33±0.58 ^a	7.43±0.45 ^{ab}	7.67±0.29 ^{ab}	9.04±0.80 ^c	1.63±0.32 ^a
C	7.32±0.56 ^{ab}	7.71±0.63 ^{ab}	10.34±0.59 ^a	5.87±0.23 ^b	7.32±0.56 ^{ab}	7.33±0.58 ^{abc}	11.94±0.82 ^a	1.60±0.10 ^a
D	7.14±0.22 ^b	7.28±0.37 ^b	10.10±0.96 ^{ab}	6.40±1.00 ^a	7.14±0.22 ^b	6.91±0.46 ^{bc}	10.45±0.50 ^b	1.37±0.37 ^b
E	8.04±0.09 ^a	8.43±0.44 ^a	9.11±0.86 ^b	6.17±0.76	8.04±0.09 ^a	7.82±0.42 ^a	8.56±0.61 ^c	1.43±0.51 ^b

All data are means of 3 triplicates expressed on dry weight. Different superscripts between columns depict significant difference ($p \leq 0.05$)

Table 10. Fungal coliforms identified cookies stored at 75% and 30% relative humidity

Samples	Fungi isolated at 70% humidity(CFU/g)				Fungi isolated at 30% humidity(CFU/g)			
	Week1	Week 3	Week 5	Week 7	Week 1	Week 3	Week 5	Week 7
A	–	–	1	1	–	–	1	1
B	–	–	1	1	–	–	1	1
C	–	–	1	1	–	–	1	1
D	–	–	1	1	–	–	1	1
E	–	–	1	1	–	–	1	1

All data are means of 3 triplicates expressed on dry weight basis. Different superscripts between columns depict significant difference ($p \leq 0.05$). A = *Saccharomyces* spp, B, C and D = *Aspergillus flavus* identified, E = *Aspergillus niger* 50 CFU = Permissible level by (Sindhu et al., 2013)

In sample B, C and D, *Aspergillus flavus* was identified. *A. flavus* are saprophytes that can be found in soil samples where they obtain their nutrition from dead and decaying matter. In particular, *A. flavus* are nuisance to farmers given that they tend to infect and contaminate crops (especially seeds). For sample E, *Aspergillus niger* was identified for cookies stored at wet and dry season conditions. *A. niger* largely exist as saprophytes, which means they obtain their nutrition from dead and decaying materials such as leaves, fruits, and other vegetables. As such they also contribute to the delay/decay of various food products, given that their source of vegetation is virtually everywhere. The fungi species isolated and identified from the cookies can be attributed to infection of the raw plant products in trace amounts. These results were lower than the 9 to 10 cfu fungi species reported for biscuit incorporated with defatted soya flour and carrot pomace powder, these values were lower than the permissible level of 50 CFU reported by Sindhu et al. [29].

4. CONCLUSION

The present results showed that, cookies can be produced from wheat, almond and carrot flour blends. Fortification of cookies with almond and carrot flour increased the vitamin content of cookies. All cookie samples were generally accepted by the sensory panelist before and after storage at all substitution levels. Cookies stored at 30% relative humidity had higher pH and lower moisture values than cookies stored at 75% relative humidity, the fungal coliforms identified in cookies were within permissible limits. Storage of cookies did not have any significant effect on the original vitamin A and C content of cookies. Variation of seasonal conditions (relative humidity) did not cause spoilage of cookies.

ACKNOWLEDGEMENTS

Thanks to the center of food technology and research (CEFTEP) Benue State University Nigeria for the study opportunity. Much thanks to Mr. Kame Stephen Mulak and Mrs. Kame Celine for financial assistance.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Baljeet SY, Ritika BY, Roshan LY. Studies on functional properties and incorporation of buckwheat flour for biscuit making. *Int. Food. Res. J.* 2010;17:1067-1076.
- Giami SY, Amusi T, Ekiyor T. Composition of bread making properties of composite flour. *Africa Journal of Biotechnology.* 2004;6(4):304-309.
- Mepha HO, Eboh SL, Nwaojigwa SU. Chemical composition, functional and baking properties of wheat plantain composite flours. *Africa Journal of Food Nutrition Deut.* 2007;7(1):1-22.
- Baljeet SY, Ritika BY, Reena K. Effect of incorporation of carrot pomace powder and germinated chickpea flour on the quality characteristics of biscuits. *International Food Research Journal.* 2014;21(1):217-222.
- Ślupski J. *Acta Sci. Pol., Technol. Aliment.* 2011;10(4):475-485.
- Dias JS. Nutritional quality and health benefits of vegetables: A review. *Food and Nutrition Sciences.* 2012;3:1354- 1374.
- Jonas T. Medicinal applications of carrot. *Journal of Food and Agric.* 2010;321:440-443.

8. Muahamad N, Faqir M.Anjum, Rai MA, Moazzam R. Khan, Shahzad Hussain, Muhammad. S. Javed. An overview of anti-nutritional factors in cereal grains with special reference to wheat-A review. Pakistan Journal of Food Sciences. 2010; 20(1-4):54-6.
9. USDA. Infant nutrition and feeding: A guide for use in WIC and CSF programs. United State Department of Agriculture Food and Nutrition Service: Special supplemental Nutrition Program for Women Infants and Children (WIC); 2009.
10. Fasanmade AA, Anyakudo MMC. Glycemic indices of selected Nigerian flour meal products in male type 2 diabetic subjects. Diabet Croat. Journal of Health Sci. 2007;36:33-38.
11. Ohimain EI. The prospects and challenges of cassava inclusion in wheat bread policy in Nigeria. International Journal of Science, Technology and Society. 2014;2: 06-17.
12. Okoye, Ebele Christiana, Onyekwelu, Chinyere Nkemakonam. Production and quality evaluation of enriched cookies from wheat, African yam bean and carrot composite flours. Annals. Food Science and Technology; 2018.
13. AOAC. Association of Official Analytical Chemists. Official methods of Analysis 18th ed. of AOAC International. Washington, DC, USA; 2011.
14. Krishnaiah D, Devi T, Bono A, Sarbatly R. Studies on phytochemical constituents of six Malaysian medicinal plants. Journal of Medicinal Plant Resources. 2009; 3(2):67-72.
15. Akinjayeju O. Quality control for the industry: A statistical approach. Lagos: Concept Publications. J. Food Science and Nutrition. 2009;4(3):364-369.
16. AOAC. Official methods of analysis. Association of official analytical chemists. 18th edition. Washington D. C. USA; 2010.
17. Association of Official Analytical Chemists. Official Methods of Analysis, 17th ed.; AOAC: Washington, DC, USA; 2003.
18. Okafor GI, Ugwu FC. Production and evaluation of cold extruded and baked read-to-eat snacks from blends of breadfruit (*Treculia africana*), cashew nut (*Anacardium occidentale*) and coconut (*cocos nucifera*). Food Science and Quality Management. 2014;23.
19. Belitz HD, Grosch W, Schieberle TY. Food Chemistry (4th Ed.). Berlin Heidelberg, Germany: Springer Verlag. 2009;403-420.
20. Baljeet SY, Ritika BY, Reena K. Effect of incorporation of carrot pomace powder and germinated chickpea flour on the quality characteristics of biscuits. International Food Research Journal. 2014;21(1):217-222.
21. Ifediba DI, Obasi CO. Evaluation of minerals, vitamins and phytochemicals in high fiber biscuits produced from blends of African Breadfruit, Maize and Coconut Flours; 2018. E-ISSN: 2454-8006.
22. Utpal Kumar Prodhana, Khan Md. Murtaja Reza Linkon, Md. Fardos Al-Amin, Md. Jahangir Alam. Development and quality evaluation of mushroom (*pleurotus sajor-caju*) enriched biscuits. Emirates Journal of Food and Agriculture. 2015;27(7):542-547.
23. Bazilla G, Rama NS, Beena MK. Physico-chemical and sensory characteristics of carrot pomace powder enriched defatted soy flour fortified biscuits. International Journal of Scientific and Research Publications. 2012;2(8). ISSN: 2250-3153.
24. William M. The Importance of Food pH in Commercial Canning Operations. FAPC-118; 2000.
25. Chinma CE, Gernah DI. Physicochemical and sensory properties of cookies produced from cassava, soybean and mango flour blends. Journal of Food Technology. 2007;5(3):256-260.
26. Lucretia I. Barber, Patience C. Obinna-Echem, Enobong M. Ogburia. Proximate composition micronutrient and sensory properties of complementary food formulated from fermented maize, soybeans and carrot flours. Sky Journal of Food Science. 2017;6(3):033–039.
27. Adeyeye, Samuel A., Akingbala, John O. Evaluation of nutritional and sensory properties of cookies produced from sweet potato- maize flour blends; 2014. ISSN: 1553-9865.
28. Omachi DO, Yusufu PA. Physico-chemical, sensory and microbiological assessment of millet based biscuits improved with

- cashew nuts (*Anarcadium occidentale*), carrot flour (*Daucus carota*). Amer. J. Food & Nutr. 2017;7(1):13-22.
29. Sindhu, Hruyial, Shweta Saloni, Harshavardhan K, Mounika B, Kalyani D, Pavankumar NS, Narayana MV. Development of biscuit incorporated with defatted soya flour and carrot pomace powder. IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT); 2013. e-ISSN: 2319-2402.

© 2020 Dinnah et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/55670>